

ON EXISTENCE AND ASYMPTOTIC BEHAVIOUR OF SOLUTIONS OF A FRACTIONAL INTEGRAL EQUATION WITH LINEAR MODIFICATION OF THE ARGUMENT

Mohamed Abdalla Darwish¹

Department of Mathematics, Massachusetts Institute of Technology
77 Massachusetts Ave., Cambridge, MA 02139-4307, USA.
E-mail: darwish@math.mit.edu

ABSTRACT. We study the solvability of a quadratic integral equation of fractional order with linear modification of the argument. This equation is considered in the Banach space of real functions defined, bounded and continuous on an unbounded interval. Moreover, we will obtain some asymptotic characterization of solutions.

AMS Mathematics Subject Classification : 45G10, 45M99, 47H09.

Keywords: Quadratic integral equation, fractional integral, existence, asymptotic behaviour, measure of noncompactness, linear modification of the argument, Schauder fixed point principal.

1. INTRODUCTION

The study of differential and integral equations with a modified argument is relatively new, it was initiated only in the past thirty years or so. These equations arise in the modeling of problems from the natural and social sciences such as biology, physics and economics, see [3, 12, 13, 14, 21, 23, 27, 29, 30, 31] and references therein. On the other hand, The first serious attempt to give a logical definition of a fractional derivative is due to Liouville, see [24] and references therein. Today differential and integral equations of fractional order play a very important role in describing numerous events and problems of the real world. For example, many problems in mechanics, physics, economics and other fields led to differential and integral equations of fractional order (cf. [9, 10, 11, 16, 17, 18, 20, 24, 32, 33]).

¹Permanent address: Department of Mathematics, Faculty of Science, Alexandria University at Damanhour, 22511 Damanhour, Egypt. E-mail: darwishma@yahoo.com.

In this paper, we will study the quadratic integral equation of fractional order with linear modification of the argument in the integral, namely

$$(1.1) \quad x(t) = a(t) + \frac{f(t, x(t))}{\Gamma(\alpha)} \int_0^t \frac{u(t, s, x(s), x(\lambda s))}{(t-s)^{1-\alpha}} ds,$$

where $t \in \mathbb{R}_+$ and $0 < \alpha, \lambda < 1$. Throughout $a : \mathbb{R}_+ \rightarrow \mathbb{R}$, $f : \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ and $u : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ are functions which satisfy special assumptions, see Section 3. Let us recall that the function $f = f(t, x)$ involves in Eq.(1.1) generates the superposition operator F defined by

$$(1.2) \quad (Fx)(t) = f(t, x(t)),$$

where $x = x(t)$ is an arbitrary function defined on \mathbb{R}_+ , see [1].

In the case $\alpha = 1$, $a(t) = 1$, $f(t, x) = x$ and $u(t, s, x, y) = \frac{t}{t+s} \phi(s) x$, where ϕ is a continuous function and $\phi(0) = 0$, Eq.(1.1) has the form

$$(1.3) \quad x(t) = 1 + x(t) \int_0^t \frac{t}{t+s} \phi(s) x(s) ds.$$

Eq.(1.3) is the Volterra counterpart of the famous quadratic integral equation of Chandrasekhar type considered in many papers and monographs (cf. [2, 6, 15, 25] for instance). Some Problems considered in the theory of radiative transfer, in the theory of neutron transport and in the kinetic theory of gases lead to quadratic integral equations (cf. [6, 8, 15, 19, 22, 25, 26, 28]).

The aim of this paper is to prove the existence of solutions of Eq.(1.1) in the space of real functions, defined, continuous and bounded on an unbounded interval. Moreover, we will obtain some asymptotic characterization of solutions of Eq.(1.1). Our proof depends on suitable combination of the technique of measures of noncompactness and the Schauder fixed point principle.

It is worthwhile mentioning that up to now the work of J. Banaś and D. O'Regan [10] is the only paper concerning with the study of quadratic integral equation of fractional order in the space of real functions defined, continuous and bounded on an unbounded interval.

2. NOTATION AND AUXILIARY FACTS

This section is devoted to collecting some definitions and results which will be needed further on. First we recall the definition of the Riemann-Liouville fractional integral [32, 33].

Definition 1. Let $f \in L_1(a, b)$, $0 \leq a < b < \infty$, and let $\alpha > 0$ be a real number. The Riemann-Liouville fractional integral of order α of the function $f(t)$ is defined by

$$I^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(s)}{(t-s)^{1-\alpha}} ds, \quad a < t < b.$$

Now, let $(E, \|\cdot\|)$ be an infinite dimensional Banach space with zero element 0. Let $B(x, r)$ denotes the closed ball centered at x with radius r . The symbol B_r stands for the ball $B(0, r)$.

If X is a subset of E , then \overline{X} and $ConvX$ denote the closure and convex closure of X , respectively. Moreover, we denote by \mathcal{M}_E the family of all nonempty and bounded subsets of E and by \mathcal{N}_E its subfamily consisting of all relatively compact subsets.

Next we give the concept of a measure of noncompactness [4]:

Definition 2. A mapping $\mu : \mathcal{M}_E \rightarrow \mathbb{R}_+ = [0, \infty)$ is said to be a measure of noncompactness in E if it satisfies the following conditions:

- 1) The family $\ker \mu = \{X \in \mathcal{M}_E : \mu(X) = 0\}$ is nonempty and $\ker \mu \subset \mathcal{N}_E$.
- 2) $X \subset Y \Rightarrow \mu(X) \leq \mu(Y)$.
- 3) $\mu(\overline{X}) = \mu(ConvX) = \mu(X)$.
- 4) $\mu(\lambda X + (1-\lambda)Y) \leq \lambda \mu(X) + (1-\lambda) \mu(Y)$ for $0 \leq \lambda \leq 1$.
- 5) If $X_n \in \mathcal{M}_E$, $X_n = \overline{X}_n$, $X_{n+1} \subset X_n$ for $n = 1, 2, 3, \dots$ and $\lim_{n \rightarrow \infty} \mu(X_n) = 0$ then $\cap_{n=1}^\infty X_n \neq \emptyset$.

The family $\ker \mu$ described above is called the kernel of the measure of noncompactness μ . Let us observe that the intersection set X_∞ from 5) belongs to $\ker \mu$. In fact, since $\mu(X_\infty) \leq \mu(X_n)$ for every n then we have that $\mu(X_\infty) = 0$.

In what follows we will work in the Banach space $BC(\mathbb{R}_+)$ consisting of all real functions defined, bounded and continuous on \mathbb{R}_+ . This space is equipped with the standard norm

$$\|x\| = \sup\{|x(t)| : t \geq 0\}.$$

Now, we recollect the construction of the measure of noncompactness in $BC(\mathbb{R}_+)$ which will be used in the next section (see [5, 7]).

Let us fix a nonempty and bounded subset X of $BC(\mathbb{R}_+)$ and numbers $\varepsilon > 0$ and $T > 0$. For arbitrary function $x \in X$ let us denote by $\omega^T(x, \varepsilon)$

the modulus of continuity of the function x on the interval $[0, T]$, i.e.

$$\omega^T(x, \varepsilon) = \sup\{|x(t) - x(s)| : t, s \in [0, T], |t - s| \leq \varepsilon\}$$

Further, let us put

$$\omega^T(X, \varepsilon) = \sup\{\omega^T(x, \varepsilon) : x \in X\},$$

$$\omega_0^T(X) = \lim_{\varepsilon \rightarrow 0} \omega^T(X, \varepsilon)$$

and

$$\omega_0^\infty(X) = \lim_{T \rightarrow \infty} \omega_0^T(X, \varepsilon).$$

Moreover, for a fixed number $t \in \mathbb{R}_+$ let us defined

$$X(t) = \{x(t) : x \in X\}$$

and

$$\text{diam}X(t) = \sup\{|x(t) - y(t)| : x, y \in X\}.$$

Finally, let us define the function μ on the family $\mathcal{M}_{BC(\mathbb{R}_+)}$ by

$$(2.1) \quad \mu(X) = \omega_0^\infty(X) + c(X),$$

where $c(X) = \limsup_{t \rightarrow \infty} \text{diam}X(t)$. The function μ is a measure of non-compactness in the space $BC(\mathbb{R}_+)$, see [5, 11]. Let us mention that the kernel $\ker \mu$ of the measure μ consists of all sets $X \in \mathcal{M}_{BC(\mathbb{R}_+)}$ such that functions from X are locally equicontinuous on \mathbb{R}_+ and vanish uniformly at infinity, i.e. for any $\varepsilon > 0$ there exists $T > 0$ such that $|x(t)| \leq \varepsilon$ for all $x \in X$ and for any $t \geq T$. This property will permit us to characterize solutions of Eq.(1.1).

3. EXISTENCE THEOREM

In this section we will study Eq.(1.1) assuming that the following hypotheses are satisfied:

- (h_1) $a : \mathbb{R}_+ \rightarrow \mathbb{R}$ is a continuous and bounded function on \mathbb{R}_+ .
- (h_2) $f : \mathbb{R}_+ \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous and there exists a continuous function $m(t) = m : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that

$$|f(t, x) - f(t, y)| \leq m(t)|x - y|$$

for all $x, y \in \mathbb{R}$ and for any $t \in \mathbb{R}_+$.

- (h_3) $u : \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous function. Moreover, there exist a function $n(t) = n : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ being continuous on \mathbb{R}_+ and a function $\Phi : \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ being continuous and nondecreasing on \mathbb{R}_+ with $\Phi(0, 0) = 0$ and such that

$$|u(t, s, x_2, y_2) - u(t, s, x_1, y_1)| \leq n(t)\Phi(|x_2 - x_1|, |y_2 - y_1|)$$

for all $t, s \in \mathbb{R}_+$ such that $t \geq s$ and for all $x_i, y_i \in \mathbb{R}$ ($i = 1, 2$).

For further purpose let us define the function $u^* : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ by $u^*(t) = \max\{|u(t, s, 0, 0)| : 0 \leq s \leq t\}$.

- (h_4) The functions $\phi, \psi, \xi, \eta : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ defined by $\phi(t) = m(t) n(t) t^\alpha$, $\psi(t) = m(t) u^*(t) t^\alpha$, $\xi = n(t)|f(t, 0)|t^\alpha$ and $\eta(t) = u^*(t)|f(t, 0)|t^\alpha$ are bounded on \mathbb{R}_+ and the functions ϕ and ξ vanish at infinity, i.e. $\lim_{t \rightarrow \infty} \phi(t) = \lim_{t \rightarrow \infty} \xi(t) = 0$.

- (h_5) There exists a positive solution r_0 of the inequality

$$(3.1) \quad \|a\| \Gamma(\alpha + 1) + [\phi^* r \Phi(r, r) + \psi^* r + \xi^* \Phi(r, r) + \eta^*] \leq r \Gamma(\alpha + 1)$$

and $\phi^* \Phi(r_0, r_0) + \psi^* < \Gamma(\alpha + 1)$, where $\phi^* = \sup\{\phi(t) : t \in \mathbb{R}_+\}$, $\psi^* = \sup\{\psi(t) : t \in \mathbb{R}_+\}$, $\xi^* = \sup\{\xi(t) : t \in \mathbb{R}_+\}$ and $\eta^* = \sup\{\eta(t) : t \in \mathbb{R}_+\}$.

Now, we are in a position to state and prove our main result.

Theorem 1. Let the hypotheses (h_1) – (h_5) be satisfied. Then Eq.(1.1) has at least one solution $x \in BC(\mathbb{R}_+)$ such that $x(t) \rightarrow 0$ as $t \rightarrow \infty$.

Proof: Denote by \mathcal{F} the operator associated with the right-hand side of equation (1.1), i.e., equation (1.1) takes the form

$$x = \mathcal{F}x,$$

where

$$(3.2) \quad (\mathcal{F}x)(t) = a(t) + (Fx)(t) \cdot (\mathcal{U}x)(t),$$

and

$$(3.3) \quad (\mathcal{U}x)(t) = \frac{1}{\Gamma(\alpha)} \int_0^t \frac{u(t, s, x(s), x(\lambda s))}{(t-s)^{1-\alpha}} ds.$$

Solving Eq.(1.1) is equivalent to finding a fixed point of the operator \mathcal{F} defined on the space $BC(\mathbb{R}_+)$.

We claim that for any function $x \in BC(\mathbb{R}_+)$ the operator \mathcal{F} is continuous on \mathbb{R}_+ . To establish this claim it suffices to show that if $x \in BC(\mathbb{R}_+)$ then $\mathcal{U}x$ is continuous function on \mathbb{R}_+ , thanks (h_1) and (h_2). For, take an arbitrary $X \in BC(\mathbb{R}_+)$ and fix $\varepsilon > 0$ and $T > 0$. Assume that

$t_1, t_2 \in \mathbb{R}_+$ are such that $|t_2 - t_1| \leq \varepsilon$. Without loss of generality we can assume that $t_2 > t_1$. Then we get

$$\begin{aligned}
|(\mathcal{U}x)(t_2) - (\mathcal{U}x)(t_1)| &= \left| \frac{1}{\Gamma(\alpha)} \int_0^{t_2} \frac{u(t_2, s, x(s), x(\lambda s))}{(t_2 - s)^{1-\alpha}} ds \right. \\
&\quad \left. - \frac{1}{\Gamma(\alpha)} \int_0^{t_1} \frac{u(t_1, s, x(s), x(\lambda s))}{(t_1 - s)^{1-\alpha}} ds \right| \\
&\leq \left| \frac{1}{\Gamma(\alpha)} \int_0^{t_2} \frac{u(t_2, s, x(s), x(\lambda s))}{(t_2 - s)^{1-\alpha}} ds \right. \\
&\quad \left. - \frac{1}{\Gamma(\alpha)} \int_0^{t_1} \frac{u(t_2, s, x(s), x(\lambda s))}{(t_2 - s)^{1-\alpha}} ds \right| \\
&\quad + \left| \frac{1}{\Gamma(\alpha)} \int_0^{t_1} \frac{u(t_2, s, x(s), x(\lambda s))}{(t_2 - s)^{1-\alpha}} ds \right. \\
&\quad \left. - \frac{1}{\Gamma(\alpha)} \int_0^{t_1} \frac{u(t_1, s, x(s), x(\lambda s))}{(t_2 - s)^{1-\alpha}} ds \right| \\
&\quad + \left| \frac{1}{\Gamma(\alpha)} \int_0^{t_1} \frac{u(t_1, s, x(s), x(\lambda s))}{(t_2 - s)^{1-\alpha}} ds \right. \\
&\quad \left. - \frac{1}{\Gamma(\alpha)} \int_0^{t_1} \frac{u(t_1, s, x(s), x(\lambda s))}{(t_1 - s)^{1-\alpha}} ds \right| \\
&\leq \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} \frac{|u(t_2, s, x(s), x(\lambda s))|}{(t_2 - s)^{1-\alpha}} ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} \frac{|u(t_2, s, x(s), x(\lambda s)) - u(t_1, s, x(s), x(\lambda s))|}{(t_2 - s)^{1-\alpha}} ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} |u(t_1, s, x(s), x(\lambda s))| [(t_1 - s)^{\alpha-1} - (t_2 - s)^{\alpha-1}] ds.
\end{aligned}$$

Therefore, if

$$\begin{aligned}
\omega_d^T(u, \varepsilon) &= \sup\{|u(t_2, s, y_1, y_2) - u(t_1, s, y_1, y_2)| : s, t_1, t_2 \in [0, T], \\
&\quad t_1 \geq s, t_2 \geq s, |t_2 - t_1| \leq \varepsilon, \text{ and } y_1, y_2 \in [-d, d]\}
\end{aligned}$$

then we obtain

$$\begin{aligned}
|(\mathcal{U}x)(t_2) - (\mathcal{U}x)(t_1)| &\leq \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} \frac{|u(t_2, s, x(s), x(\lambda s)) - u(t_2, s, 0, 0)| + |u(t_2, s, 0, 0)|}{(t_2 - s)^{1-\alpha}} ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} \frac{\omega_{\|x\|}^T(u, \varepsilon)}{(t_2 - s)^{1-\alpha}} ds \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} [|u(t_1, s, x(s), x(\lambda s)) - u(t_1, s, 0, 0)| + |u(t_1, s, 0, 0)|] \\
&\quad \quad \times [(t_1 - s)^{\alpha-1} - (t_2 - s)^{\alpha-1}] ds \\
&\leq \frac{1}{\Gamma(\alpha)} \int_{t_1}^{t_2} \frac{n(t_2)\Phi(|x(s)|, |x(\lambda s)|) + u^*(t_2)}{(t_2 - s)^{1-\alpha}} ds \\
&\quad + \frac{\omega_{\|x\|}^T(u, \varepsilon)}{\Gamma(\alpha + 1)} [t_2^\alpha - (t_2 - t_1)^\alpha] \\
&\quad + \frac{1}{\Gamma(\alpha)} \int_0^{t_1} [n(t_1)\Phi(|x(s)|, |x(\lambda s)|) + u^*(t_1)] \\
&\quad \quad \times [(t_1 - s)^{\alpha-1} - (t_2 - s)^{\alpha-1}] ds \\
&\leq \frac{n(t_2)\Phi(\|x\|, \|x\|) + u^*(t_2)}{\Gamma(\alpha + 1)} (t_2 - t_1)^\alpha + \frac{\omega_{\|x\|}^T(u, \varepsilon)}{\Gamma(\alpha + 1)} t_1^\alpha \\
&\quad + \frac{n(t_1)\Phi(\|x\|, \|x\|) + u^*(t_1)}{\Gamma(\alpha + 1)} [t_1^\alpha - t_2^\alpha + (t_2 - t_1)^\alpha].
\end{aligned}$$

Thus

(3.4)

$$\omega^T(\mathcal{U}x, \varepsilon) \leq \frac{1}{\Gamma(\alpha + 1)} \{2\varepsilon^\alpha [\hat{n}(T)\Phi(\|x\|, \|x\|) + \hat{u}(T)] + T^\alpha \omega_{\|x\|}^T(u, \varepsilon)\},$$

where we denoted

$$\hat{n}(T) = \max\{n(t) : t \in [0, T]\}$$

and

$$\hat{u}(T) = \max\{u^*(t) : t \in [0, T]\}.$$

In view of the uniform continuity of the function u on $[0, T] \times [0, T] \times [-\|x\|, \|x\|] \times [-\|x\|, \|x\|]$ we have that $\omega_{\|x\|}^T(u, \varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. From the above inequality we infer that the function $\mathcal{U}x$ is continuous on the interval $[0, T]$ for any $T > 0$. This yields the continuity of $\mathcal{U}x$ on \mathbb{R}_+ , and consequently, the function $\mathcal{F}x$ is continuous on \mathbb{R}_+ .

Now, we show that $\mathcal{F}x$ is bounded on \mathbb{R}_+ . Indeed, in view of our hypotheses for arbitrary $x \in BC(\mathbb{R}_+)$ and for a fixed $t \in \mathbb{R}_+$ we have

$$\begin{aligned}
|(\mathcal{F}x)(t)| &\leq \left| a(t) + \frac{f(t, x(t))}{\Gamma(\alpha)} \int_0^t \frac{u(t, s, x(s), x(\lambda s))}{(t-s)^{1-\alpha}} ds \right| \\
&\leq \|a\| + \frac{1}{\Gamma(\alpha)} [|f(t, x(t)) - f(t, 0)| + |f(t, 0)|] \\
&\quad \times \int_0^t \frac{|u(t, s, x(s), x(\lambda s)) - u(t, s, 0, 0)| + |u(t, s, 0, 0)|}{(t-s)^{1-\alpha}} ds \\
&\leq \|a\| + \frac{m(t)\|x\| + |f(t, 0)|}{\Gamma(\alpha)} \int_0^t \frac{n(t)\Phi(|x(s)|, |x(\lambda s)|) + u^*(t)}{(t-s)^{1-\alpha}} ds \\
&\leq \|a\| + \frac{m(t)\|x\| + |f(t, 0)|}{\Gamma(\alpha+1)} [n(t)\Phi(\|x\|, \|x\|) + u^*(t)] t^\alpha \\
&= \|a\| + \frac{1}{\Gamma(\alpha+1)} [\phi(t)\|x\|\Phi(\|x\|, \|x\|) + \psi(t)\|x\| + \xi(t)\Phi(\|x\|, \|x\|) + \eta(t)].
\end{aligned}$$

Hence, $\mathcal{F}x$ is bounded on \mathbb{R}_+ , thanks hypothesis (h_4) . This assertion in conjunction with the continuity of $\mathcal{F}x$ on \mathbb{R}_+ allows us to conclude that the operator \mathcal{F} maps $BC(\mathbb{R}_+)$ into itself. Moreover, from the last estimate we have

$$\|\mathcal{F}x\| \leq \|f\| + \frac{1}{\Gamma(\alpha+1)} [\phi^*\|x\|\Phi(\|x\|, \|x\|) + \psi^*\|x\| + \xi^*\Phi(\|x\|, \|x\|) + \eta^*].$$

Linking this estimate with hypothesis (h_5) we deduce that there exists $r_0 > 0$ such that the operator \mathcal{F} transforms the ball B_{r_0} into itself.

Now, we prove that the operator \mathcal{F} is continuous on the ball B_{r_0} . To do this, let us fix $\varepsilon > 0$ and take $x, y \in B_{r_0}$ such that $\|x - y\| \leq \varepsilon$. Then,

for $t \in \mathbb{R}_+$ we get

$$\begin{aligned}
|(\mathcal{F}x)(t) - (\mathcal{F}y)(t)| &\leq \left| \frac{f(t, x(t))}{\Gamma(\alpha)} \int_0^t \frac{u(t, s, x(s), x(\lambda s))}{(t-s)^{1-\alpha}} ds \right. \\
&\quad \left. - \frac{f(t, y(t))}{\Gamma(\alpha)} \int_0^t \frac{u(t, s, y(s), y(\lambda s))}{(t-s)^{1-\alpha}} ds \right| \\
&\leq \frac{|f(t, x(t)) - f(t, y(t))|}{\Gamma(\alpha)} \int_0^t \frac{|u(t, s, x(s), x(\lambda s))|}{(t-s)^{1-\alpha}} ds \\
&\quad + \frac{|f(t, y(t))|}{\Gamma(\alpha)} \int_0^t \frac{|u(t, s, x(s), x(\lambda s)) - u(t, s, y(s), y(\lambda s))|}{(t-s)^{1-\alpha}} ds \\
&\leq \frac{m(t)|x(t) - y(t)|}{\Gamma(\alpha)} \int_0^t \frac{|u(t, s, x(s), x(\lambda s)) - u(t, s, 0, 0)| + |u(t, s, 0, 0)|}{(t-s)^{1-\alpha}} ds \\
&\quad + \frac{m(t)|y(t)| + |f(t, 0)|}{\Gamma(\alpha)} \int_0^t \frac{n(t) \Phi(|x(s) - y(s)|, |x(\lambda s) - y(\lambda s)|)}{(t-s)^{1-\alpha}} ds \\
&\leq \frac{m(t)|x(t) - y(t)|}{\Gamma(\alpha)} \int_0^t \frac{n(t) \Phi(|x(s)|, |x(\lambda s)|) + u^*(t)}{(t-s)^{1-\alpha}} ds \\
&\quad + \frac{m(t)|y(t)| + |f(t, 0)|}{\Gamma(\alpha)} \int_0^t \frac{n(t) \Phi(|x(s)| + |y(s)|, |x(\lambda s)| + |y(\lambda s)|)}{(t-s)^{1-\alpha}} ds \\
&\leq \frac{|x(t) - y(t)|}{\Gamma(\alpha + 1)} [\phi(t) \Phi(\|x\|, \|x\|) + \psi(t)] \\
&\quad + \frac{\phi(t)|y(t)| \Phi(\|x\| + \|y\|, \|x\| + \|y\|)}{\Gamma(\alpha + 1)} \\
&\quad + \frac{\xi(t) \Phi(\|x\| + \|y\|, \|x\| + \|y\|)}{\Gamma(\alpha + 1)} \\
&\leq \frac{1}{\Gamma(\alpha + 1)} [\varepsilon \phi(t) \Phi(r_0, r_0) + \varepsilon \psi(t) \\
&\quad + \phi(t) r_0 \Phi(2r_0, 2r_0) + \xi(t) \Phi(2r_0, 2r_0)].
\end{aligned} \tag{3.5}$$

Thus \mathcal{F} is continuous on B_{r_0} , thanks hypothesis (h_4) .

In what follows let us take a nonempty set $X \subset B_{r_0}$. Then, for arbitrary $x, y \in X$ and for a fixed $t \in \mathbb{R}_+$, calculating in the same way as in

estimate (3.5), we obtain

$$\begin{aligned}
|(\mathcal{F}x)(t) - (\mathcal{F}y)(t)| &\leq \frac{|x(t) - y(t)|}{\Gamma(\alpha + 1)} [\phi(t)\Phi(\|x\|, \|x\|) + \psi(t)] \\
&\quad + \frac{\Phi(\|x\| + \|y\|, \|x\| + \|y\|)}{\Gamma(\alpha + 1)} [\phi(t)\|y\| + \xi(t)] \\
&\leq \frac{\phi(t)\Phi(r_0, r_0) + \psi(t)}{\Gamma(\alpha + 1)} |x(t) - y(t)| \\
&\quad + \frac{\Phi(2r_0, 2r_0)}{\Gamma(\alpha + 1)} [\phi(t)r_0 + \xi(t)].
\end{aligned}$$

Hence, we can easily deduce the following inequality

$$\text{diam}(\mathcal{F}X)(t) \leq \frac{\phi(t)\Phi(r_0, r_0) + \psi(t)}{\Gamma(\alpha + 1)} \text{diam}X(t) + \frac{\Phi(2r_0, 2r_0)}{\Gamma(\alpha + 1)} [\phi(t)r_0 + \xi(t)].$$

Now, taking into account hypothesis (h_4) we obtain

$$(3.6) \quad c(\mathcal{F}X) \leq k c(X),$$

where we denoted $k = \frac{\phi^*\Phi(r_0, r_0) + \psi^*}{\Gamma(\alpha + 1)}$. Obviously, in view of hypothesis (h_5) we have that $k < 1$.

In what follows, let us take arbitrary numbers $\varepsilon > 0$ and $T > 0$. Choose a function $x \in X$ and take $t_1, t_2 \in [0, T]$ such that $|t_2 - t_1| \leq \varepsilon$. Without loss of generality we can assume that $t_2 > t_1$. Then, taking into

account our hypotheses and (3.4), we have

$$\begin{aligned}
|(\mathcal{F}x)(t_2) - (\mathcal{F}x)(t_1)| &\leq |a(t_2) - a(t_1)| + |(Fx)(t_2)(\mathcal{U}x)(t_2) - (Fx)(t_1)(\mathcal{U}x)(t_2)| \\
&\quad + |(Fx)(t_1)(\mathcal{U}x)(t_2) - (Fx)(t_1)(\mathcal{U}x)(t_1)| \\
&\leq \omega^T(a, \varepsilon) + \frac{|f(t_2, x(t_2)) - f(t_1, x(t_1))|}{\Gamma(\alpha)} \\
&\quad \times \int_0^{t_2} \frac{|u(t_2, s, x(s), x(\lambda s)) - u(t_2, s, 0, 0)| + |u(t_2, s, 0, 0)|}{(t_2 - s)^{1-\alpha}} ds \\
&\quad + \frac{|f(t_1, x(t_1)) - f(t_1, 0)| + |f(t_1, 0)|}{\Gamma(\alpha + 1)} \left\{ 2\varepsilon^\alpha [\hat{n}(T)\Phi(\|x\|, \|x\|) + \hat{u}(T)] + T^\alpha \omega_{\|x\|}^T(u, \varepsilon) \right\} \\
&\leq \omega^T(a, \varepsilon) + \frac{m(t_2)|x(t_2) - x(t_1)| + \omega_f^T(\varepsilon)}{\Gamma(\alpha)} \int_0^{t_2} \frac{n(t_2)\Phi(|x(s)|, |x(\lambda s)|) + u^*(t_2)}{(t_2 - s)^{1-\alpha}} ds \\
&\quad + \frac{m(t_1)|x(t_1)| + |f(t_1, 0)|}{\Gamma(\alpha + 1)} \left\{ 2\varepsilon^\alpha [\hat{n}(T)\Phi(\|x\|, \|x\|) + \hat{u}(T)] + T^\alpha \omega_{\|x\|}^T(u, \varepsilon) \right\} \\
&\leq \omega^T(a, \varepsilon) + \frac{t_2^\alpha}{\Gamma(\alpha + 1)} [m(t_2)\omega^T(x, \varepsilon) + \omega_f^T(\varepsilon)] [n(t_2)\Phi(r_0, r_0) + u^*(t_2)] \\
&\quad + \frac{\hat{m}(T)r_0 + \hat{f}(T)}{\Gamma(\alpha + 1)} \left\{ 2\varepsilon^\alpha [\hat{n}(T)\Phi(\|x\|, \|x\|) + \hat{u}(T)] + T^\alpha \omega_{\|x\|}^T(u, \varepsilon) \right\} \\
&\leq \omega^T(a, \varepsilon) + \frac{[\phi(t_2)\Phi(r_0, r_0) + \psi(t_2)]}{\Gamma(\alpha + 1)} \omega^T(x, \varepsilon) + \frac{T^\alpha \omega_f^T(\varepsilon)}{\Gamma(\alpha + 1)} [\hat{n}(T)\Phi(r_0, r_0) + \hat{u}(T)] \\
&\quad + \frac{\hat{m}(T)r_0 + \hat{f}(T)}{\Gamma(\alpha + 1)} \left\{ 2\varepsilon^\alpha [\hat{n}(T)\Phi(\|x\|, \|x\|) + \hat{u}(T)] + T^\alpha \omega_{\|x\|}^T(u, \varepsilon) \right\} \\
&\leq \omega^T(a, \varepsilon) + \frac{[\phi^*\Phi(r_0, r_0) + \psi^*]}{\Gamma(\alpha + 1)} \omega^T(x, \varepsilon) + \frac{T^\alpha \omega_f^T(\varepsilon)}{\Gamma(\alpha + 1)} [\hat{n}(T)\Phi(r_0, r_0) + \hat{u}(T)] \\
&\quad (3.7) + \frac{\hat{m}(T)r_0 + \hat{f}(T)}{\Gamma(\alpha + 1)} \left\{ 2\varepsilon^\alpha [\hat{n}(T)\Phi(\|x\|, \|x\|) + \hat{u}(T)] + T^\alpha \omega_{\|x\|}^T(u, \varepsilon) \right\},
\end{aligned}$$

where we denoted

$$\omega_f^T(\varepsilon) = \sup\{|f(t_2, x(t_2)) - f(t_1, x(t_1))| : t_1, t_2 \in [0, T], |t_2 - t_1| \leq \varepsilon, x \in [-r_0, r_0]\},$$

$$\hat{m}(T) = \max\{m(t) : t \in [0, T]\}$$

and

$$\hat{f}(T) = \max\{|f(t, 0)| : t \in [0, T]\}.$$

Now, keeping in mind the uniform continuity of the function $f = f(t, x)$ on the set $[0, T] \times [r_0, r_0]$ and the uniform continuity of the function $u = u(t, s, x, y)$ on the set $[0, T] \times [0, T] \times [r_0, r_0] \times [r_0, r_0]$, from estimate

(3.7) we derive the following one

$$\omega_0^T(\mathcal{F}X) \leq k \omega_0^T(X).$$

Hence we have

$$(3.8) \quad \omega_0^\infty(\mathcal{F}X) \leq k \omega_0^\infty(X).$$

From (3.6) and (3.8) and the definition of the measure of noncompactness μ given by formula (2.1), we obtain

$$(3.9) \quad \mu(\mathcal{F}X) \leq k \mu(X).$$

In the sequel let us put $B_{r_0}^1 = \text{Conv}\mathcal{F}(B_{r_0})$, $B_{r_0}^2 = \text{Conv}\mathcal{F}(B_{r_0}^1)$ and so on. In this way we have constructed a decreasing sequence of nonempty, bounded, closed and convex subsets $(B_{r_0}^n)$ of B_{r_0} such that $\mathcal{F}(B_{r_0}^n) \subset B_{r_0}^n$ for $n = 1, 2, \dots$. Since the above reasons leading to (3.9) holds for any subset X of B_{r_0} we have

$$\mu(B_{r_0}^n) \leq k^n \mu(B_{r_0}), \text{ for any } n = 1, 2, 3, \dots$$

This implies that $\lim_{n \rightarrow \infty} \mu(B_{r_0}^n) = 0$. Hence, taking into account Definition 2 we infer that the set $Y = \bigcap_{n=1}^{\infty} B_{r_0}^n$ is nonempty, bounded, closed and convex subset of B_{r_0} . Moreover, $Y \in \ker \mu$. Also, the operator \mathcal{F} maps Y into itself.

We will prove that the operator \mathcal{F} is continuous on the set Y . In order to do this let us fix a number $\varepsilon > 0$ and take arbitrary functions $x, y \in Y$ such that $\|x - y\| \leq \varepsilon$. Keeping in mind the facts that $Y \in \ker \mu$ and the structure of sets belong to $\ker \mu$ we can find a number $T > 0$ such that for each $z \in Y$ and $t \geq T$ we have that $|z(t)| \leq \varepsilon$. Since \mathcal{F} maps Y into itself we have that $\mathcal{F}x, \mathcal{F}y \in Y$. Thus, for $t \geq T$ we get

$$(3.10) \quad |(\mathcal{F}x)(t) - (\mathcal{F}y)(t)| \leq |(\mathcal{F}x)(t)| + |(\mathcal{F}y)(t)| \leq 2\varepsilon.$$

On the other hand, let us assume $t \in [0, T]$. Then we obtain

$$\begin{aligned}
|(\mathcal{F}x)(t) - (\mathcal{F}y)(t)| &\leq \frac{m(t)|x(t) - y(t)|}{\Gamma(\alpha)} \int_0^t \frac{n(s) \Phi(|x(s)|, |x(\lambda s)|) + u^*(t)}{(t-s)^{1-\alpha}} ds \\
&\quad + \frac{m(t)|y(t)| + |f(t, 0)|}{\Gamma(\alpha)} \int_0^t \frac{n(s) \Phi(|x(s) - y(s)|, |x(\lambda s) - y(\lambda s)|)}{(t-s)^{1-\alpha}} ds \\
&\leq \frac{[m(t)n(t)\Phi(r_0, r_0) + m(t)u^*(t)]\varepsilon}{\Gamma(\alpha)} \int_0^t \frac{ds}{(t-s)^{1-\alpha}} \\
&\quad + \frac{[m(t)n(t)r_0 + n(t)|f(t, 0)]\Phi(\varepsilon, \varepsilon)}{\Gamma(\alpha)} \int_0^t \frac{ds}{(t-s)^{1-\alpha}} \\
&\leq \frac{\phi(t)\Phi(r_0, r_0) + \psi(t)}{\Gamma(\alpha+1)} \varepsilon + \frac{\phi(t)r_0 + \xi(t)}{\Gamma(\alpha+1)} \Phi(\varepsilon, \varepsilon) \\
(3.11) \quad &\leq \frac{\phi^*\Phi(r_0, r_0) + \psi^*}{\Gamma(\alpha+1)} \varepsilon + \frac{\phi^*r_0 + \xi^*}{\Gamma(\alpha+1)} \Phi(\varepsilon, \varepsilon).
\end{aligned}$$

Now, taking into account (3.10), (3.11) and hypothesis (h_4) we conclude that the operator \mathcal{F} is continuous on the set Y .

Finally, linking all above obtained facts about the set Y and the operator $\mathcal{F} : Y \rightarrow Y$ and using the classical Schauder fixed point principal we deduce that the operator \mathcal{F} has at least one fixed point x in the set Y . Obviously the function $x = x(t)$ is a solution of the quadratic integral equation (1.1). Moreover, since $Y \in \ker \mu$ we have that $x(t) \rightarrow 0$ as $t \rightarrow \infty$. This completes the proof.

4. EXAMPLE

Consider the following quadratic integral equation of fractional order with linear modification of the argument, $\alpha = \frac{1}{2}$,

$$(4.12) \quad x(t) = te^{-t} + \frac{t + t^2x(t)}{2\Gamma(\frac{1}{2})} \int_0^t \frac{(|x(s)| + |x(\lambda s)|)e^{-2t-s} + 1/(1 + 5t^{5/2})}{\sqrt{t-s}} ds.$$

In this example, we have that $a(t) = te^{-t}$ and this function satisfies hypothesis (h_1) and $\|a\| = a(1) = 1/e$. Moreover, $f(t, x(t)) = (t + t^2x(t))/2$ and satisfies hypothesis (h_2) with $m(t) = t^2/2$, and $|f(t, 0)| = f(t, 0) = t/2$. Also, $u(t, s, x, y) = (x + y)e^{-2t-s} + 1/(1 + 5t^{5/2})$ verifies hypothesis (h_3) with $n(t) = e^{-2t}$, $\phi(x, y) = x + y$ and $u(t, s, 0, 0) = 1/(1 + 5t^{5/2})$. Now, we have

$$\phi(t) = \frac{1}{2}t^{5/2}e^{-2t},$$

$$\psi(t) = \frac{t^{5/2}}{2(1 + 5t^{5/2})},$$

$$\xi(t) = \frac{1}{2}t^{3/2}e^{-2t}$$

and

$$\eta(t) = \frac{t^{3/2}}{2(1 + 5t^{5/2})}.$$

It is easy to see that the functions ϕ , ψ , ξ and η are bounded on \mathbb{R}_+ and also $\lim_{t \rightarrow \infty} \phi(t) = \lim_{t \rightarrow \infty} \xi(t) = 0$. Hence, hypothesis (h_4) is satisfied. Moreover, we have

$$\phi^* = \phi(5/4) = \frac{1}{2}(5/4)^{5/2}e^{-5/2} = 0.0716982...,$$

$$\psi^* = 0.1,$$

$$\xi^* = \phi(3/4) = \frac{1}{2}(3/4)^{5/2}e^{-3/2} = 0.0543477...$$

and

$$\eta^* = \eta((0.2)^{2/5}) = 0.0410503... .$$

In this case the inequality (3.1) has the form

$$(4.13) \quad \Gamma\left(\frac{3}{2}\right)e^{-1} + 2r^2\phi^* + r\psi^* + 2r\xi^* + \eta^* \leq r\Gamma\left(\frac{3}{2}\right).$$

Let us denote by $H(r)$ the left hand side of the last inequality, i.e.

$$H(r) = \Gamma\left(\frac{3}{2}\right)e^{-1} + 2r^2\phi^* + r\psi^* + 2r\xi^* + \eta^*.$$

For $r = 1$ we obtain

$$\begin{aligned} H(1) &= \Gamma\left(\frac{3}{2}\right)e^{-1} + 2\phi^* + \psi^* + 2\xi^* + \eta^* \\ &= \Gamma\left(\frac{3}{2}\right) 0.3678794... + 0.3931423... . \end{aligned}$$

Hence, inequality (4.13) admits $r_0 = 1$ as a positive solution since $\Gamma(\frac{3}{2}) \simeq 0.886227$. Moreover,

$$\phi^*\phi(r_0, r_0) + \psi^* \simeq 0.2433966 < \Gamma\left(\frac{3}{2}\right).$$

Therefore, Theorem 1 guarantees that equation (4.12) has a solution $x = x(t)$ in the space \mathbb{R}_+ belonging to the ball B_1 such that $x(t) \rightarrow 0$ as $t \rightarrow \infty$.

REFERENCES

- [1] J. APPELL AND P.P. ZABREJKO, *Nonlinear superposition Operators*, Cambridge tracts in mathematics Vol.**95**, Cambridge University Press, 1990.
- [2] I.K. ARGYROS, *Quadratic equations and applications to Chandrasekhar's and related equations*, Bull. Austral. Math. Soc. **32** (1985), 275-292.
- [3] D. BAINOV AND D. MISHEV, *Oscillation Theory for Neutral Differential Equations with Delay*, Hilger, Bristol, 1991.
- [4] J. BANAŚ AND K. GOEBEL, *Measures of noncompactness in Banach spaces*, Lecture Notes in Pure and Applied Mathematics **60**, Marcel Dekker, New York, 1980.
- [5] J. BANAŚ, *Measures of noncompactness in the space of continuous tempered functions*, Demonstratio Math. **14** (1981), 127-133.
- [6] J. BANAŚ, M. LECKO AND W.G. EL-SAYED, *Existence theorems of some quadratic integral equations*, J. Math. Anal. Appl. **222** (1998), 276-285.
- [7] J. BANAŚ AND L. OLSZOWY, *Measures of noncompactness related to monotonicity*, Comment. Math. **41** (2001), 13-23.
- [8] J. BANAŚ AND B. RZEPKA, *On existence and asymptotic stability of solutions of a nonlinear integral equation*, J. Math. Anal. Appl. **284** (2003), 165-173.
- [9] J. BANAŚ AND B. RZEPKA, *Monotonic solutions of a quadratic integral equation of fractional order*, J. Math. Anal. Appl. **332** (2007), 1370-1378.
- [10] J. BANAŚ AND D. O'REGAN, *On existence and local attractivity of solutions of a quadratic integral equation of fractional order*, J. Math. Anal. Appl. (2008), doi: 10.1016/j.jmaa.2008-04-050.
- [11] J. BANAŚ AND L. OLSZOWY, *On a class of measures of noncompactness in Banach algebras and their application to nonlinear integral equations*, Reprint.
- [12] M. BENCHOHRA AND M.A. DARWISH *On Unique Solvability of Quadratic Integral Equations with Linear Modification of the Argument*, Accepted for publication in Miskolc Math. Notes (2008).
- [13] J. Caballero, B. López and K. Sadarangani, Existence of nondecreasing and continuous solutions of an integral equation with linear modification of the argument, *Acta Math. Sin. (Engl. Ser.)* **2** (9) (2007), 1719-1728.
- [14] J. Carr and J. Dyson, The functional differential equation $y'(x) = ay(\lambda x) + by(x)$, *Proc. Roy. Soc. Edinburgh Sect. A* **74** (1974/75) (1976), 165-174.
- [15] S. CHANDRASEKHAR, *Radiative Transfer*, Oxford University Press, London, 1950.
- [16] M.A. DARWISH, *On monotonic solutions of an integral equation of Abel type*, Accepted for publication in Mathematica Bohemica.
- [17] M.A. DARWISH, *On monotonic solutions of a singular quadratic integral equation with supremum*, Dynam. Syst. Appl. (2008).
- [18] M.A. DARWISH, *On solvability of some quadratic functional-integral equation in Banach algebra*, Commun. Appl. Anal. **11** (3-4)(2007), 441-450.
- [19] M.A. DARWISH, *On global attractivity of solutions of a functional-integral equation*, Electron. J. Qual. Theory Differ. Equ. No. 21 (2007) 10pp.
- [20] M.A. DARWISH, *On quadratic integral equation of fractional orders*, J. Math. Anal. Appl. **311** (2005), 112-119.

- [21] M.A. DARWISH, *On a quadratic fractional integral equation with linear modification of the argument*, Reprint.
- [22] K. DEIMLING, *Nonlinear functional analysis*, Springer-Verlag, Berlin, 1985.
- [23] G.M. Dunkel, *Function Differential Equations: Examples and Problems*, Lecture Notes in Math. **144**, Springer, Berlin, 1970.
- [24] R. HILFER, *Applications of Fractional calculus in Physics*, World Scientific, Singapore, 2000.
- [25] S. HU, M. KHAVANI AND W. ZHUANG, *Integral equations arising in the kinetic theory of gases*, Appl. Analysis, **34** (1989), 261-266.
- [26] C.T. KELLEY, *Approximation of solutions of some quadratic integral equations in transport theory*, J. Integral Eq. **4** (1982), 221-237.
- [27] M.R.S. KULENOVIĆ, *Oscillation of the Euler differential equation with delay*, Czechoslovak Math. J. **45 (120)** (1) (1995), 1-6.
- [28] R.W. LEGGETT, *A new approach to the H-equation of Chandrasekher*, SIAM J. Math. **7** (1976), 542-550.
- [29] H. MELVIN, *A family of solutions of the IVP for the equation $x'(t) = ax(\lambda t)$, $\lambda > 1$* , Aequationes Math. **9** (1973), 273-280.
- [30] V. MUREŞAN, *Volterra integral equations with iterations of linear modification of the argument*, Novi Sad J. Math. **33** (2) (2003), 1-10.
- [31] V. MUREŞAN, *On a class of Volterra integral equations with deviating argument*, Studia Univ. Babeş-Bolyai Math. **44** (1) (1999), 47-54.
- [32] I. PODLUBNY, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [33] S.G. SAMKO, A.A. KILBAS AND O.I. MARICHEV, *Fractional integrals and derivatives: theory and applications*, Gordon and Breach Science Publ., Amsterdam, 1993. [Russian Edition 1987]